


# A large handaxe from Wadi Dabsa and early hominin adaptations within the Arabian Peninsula

Frederick W.F. Foulds<sup>1,\*</sup>, Andrew Shuttleworth<sup>2</sup>, Anthony Sinclair<sup>3</sup>, Abdullah M. Alsharekh<sup>4</sup>, Saud Al Ghamdi<sup>5</sup>, Robyn H. Inglis<sup>6,7</sup> & Geoff N. Bailey<sup>6</sup>



*The role played by the Arabian Peninsula in hominin dispersals out of Africa has long been debated. The DISPERSE Project has focused on south-western Arabia as a possible centre of hominin settlement and a primary stepping-stone for such dispersals. This work has led to the recent discovery, at Wadi Dabsa, of an exceptional assemblage of over 1000 lithic artefacts, including the first known giant handaxe from the Arabian Peninsula. The site and its associated artefacts provide important new evidence for hominin dispersals out of Africa, and give further insight into the giant handaxe phenomenon present within the Acheulean stone tool industry.*

**Keywords:** Saudi Arabia, Palaeolithic, Acheulean, human evolution, handaxes, hominin dispersal

## Introduction

Acheulean bifacial tools are considered to be one of the greatest enigmas of the African Early Stone Age and European Lower Palaeolithic (Wymer 1982: 102). They appear in

<sup>1</sup> Department of Archaeology, Durham University, South Road, Durham DH1 3LE, UK

<sup>2</sup> Department of Anthropology, Durham University, South Road, Durham DH1 3LE, UK

<sup>3</sup> Department of Archaeology, Classics and Egyptology, University of Liverpool, 12–14 Abercromby Square, Liverpool L69 7WZ, UK

<sup>4</sup> Department of Archaeology, King Saud University, P.O. Box 2627, Riyadh 12372, Kingdom of Saudi Arabia

<sup>5</sup> Qatar Museums Authority, QM Tower, P.O. Box 2777, Doha, State of Qatar

<sup>6</sup> Department of Archaeology, University of York, King's Manor, York YO1 7EP, UK

<sup>7</sup> Department of Environmental Sciences, Macquarie University, Sydney, NSW 2109, Australia

\* Author for correspondence (Email: [frederick.foulds@googlemail.com](mailto:frederick.foulds@googlemail.com))

the archaeological record from 1.76 million years ago and persist for over one million years, representing an extensive period of technological stasis associated with a variety of hominin species, landscapes and environments (Asfaw *et al.* 1992; Quade *et al.* 2004; Lepre *et al.* 2011). Wrapped within this technological enigma are overly large handaxes, whose excesses in both size and weight have confounded archaeologists as to the exact purpose behind their production (Wynn 1995). Here we present the recently discovered Palaeolithic site of Wadi Dabsa, Saudi Arabia, and the recovery of a large Acheulean handaxe. The rich lithic assemblage from Wadi Dabsa not only yields evidence of how hominin populations may have adapted to varied landscapes and conditions during their dispersals out of Africa, but also provides insight into how such large bifacial tools may have been used.

There is clear evidence that the Arabian Peninsula was host to Acheulean hominins throughout the Early to Middle Pleistocene (*c.* 2 Mya–200 kya), and that these hominins occupied landscapes and environments close to water and raw material sources in the interior and coastal regions of the Peninsula (Petraglia 2003; Field & Lahr 2005; Bailey *et al.* 2007, 2015; Petraglia & Rose 2009; Petraglia *et al.* 2009; Groucutt & Petraglia 2012). It is not clear whether a land bridge would have existed across the southern end of the Red Sea at periods of low sea level during the Pliocene or Early Pleistocene. The long-term rotation of the Arabian Plate away from Africa might imply progressive widening of the sea channel and therefore the possible existence of a land bridge at some earlier time. Accommodation of plate motions by crustal deformation, however, mainly occurs in the Afar depression and along the Arabian escarpment, rather than in the area of the Red Sea Channel; there are too many uncertainties concerning the topographic impact of tectonic and volcanic activity at this early period to be certain. Nonetheless, by the Middle Pleistocene, and certainly from approximately half a million years ago, analysis of isotopic composition in deep sea cores and from tectonic modelling of palaeocoastlines shows that a narrow and shallow sea connection to the Indian Ocean persisted for long periods during lower sea levels in the Hanish Sill region. This would have afforded the possibility of sea crossings of no more than 4km. A very extensive area of potentially attractive coastal lowland would also have been exposed on both sides of this channel (Siddall *et al.* 2003; Bailey 2009; Lambeck *et al.* 2011; Rohling *et al.* 2013; Bailey *et al.* 2015).

The significance of Arabia in the dispersal and evolution of hominins out of Africa is, however, much debated, due to the lack of chronological certainty for many of its prehistoric sites; and although the use of the ‘Southern Dispersal Route’ (involving a crossing of the southern Red Sea and the southern Arabian Peninsula) during the Pleistocene is plausible, the lack of significant genetic input from within modern populations in Arabia suggests that these migrations involved small populations (Cabrera *et al.* 2009). Regardless of their size, these groups would have migrated into, and along, what are now the Red Sea and Gulf of Aden coastlines. These regions, in particular those along the southern Red Sea coast with their added increment of territory available at lower sea levels, would have presented hominins with a productive landscape of fauna, water and raw material sources comparable to those already experienced in the Horn of Africa. They would also have acted as refugia during periods of hyperaridity, when the Arabian interior would have become uninhabitable (Petraglia & Rose 2009; Winder *et al.* 2015).

## **The DISPERSE Project and Wadi Dabsa**

The DISPERSE Project is concerned with the impact of sea-level change and active tectonics on the early landscapes of human evolution and hominin dispersal within Africa and beyond (Bailey *et al.* 2012, 2015; Devès *et al.* 2014; Inglis *et al.* 2014a & b; Kübler *et al.* 2016). Work has concentrated in particular on the southern Red Sea and the south-western Arabian escarpment; on reconstruction of prehistoric landscapes on land and underwater; and on survey and investigation of Palaeolithic sites and later coastal middens in their landscape setting. This regional focus is informed by the hypothesis that south-western Arabia was an early centre of hominin settlement and a primary stepping-stone for range expansion out of Africa. This hypothesis is based on the presence of similar tectonic and volcanic landscapes that were advantageous in the earliest centres of human evolution in the East African Rift, proximity and accessibility to the Rift across a narrow sea-crossing for long periods of the Pleistocene and on relatively beneficial climatic conditions and ecological diversity (King & Bailey 2006; Bailey *et al.* 2007, 2012, 2015; Reynolds *et al.* 2011; Winder *et al.* 2013, 2015).

The Harrat Al Birk is an extensive series of basaltic flows that extend along the present-day coastline for approximately 100km, and stretch inland for around 30km, where they meet the basement rocks of the foothills of the Western Arabian Escarpment (Dabbagh *et al.* 1984; Prinz 1984). Wadi Dabsa, at present a seasonally flowing watercourse running for approximately 7km to the sea, drains the western edge of the Harrat (Figures 1 & 2). In its upper reaches, the wadi flows through a small basin within the basalt, the base of which has been almost completely covered by tufa deposition, around 1km<sup>2</sup> in total. The tufa was deposited during a past period of consistent flow of carbonate-rich water, possibly fed by a number of small tributaries draining the surrounding slopes, and forming a series of dams and pools (Inglis *et al.* 2015). The tufa formation suggests perennial water flow, and, given the limited catchment of the basin, may be linked to past spring activity, rather than runoff. Regardless of the source, the presence of large volumes of water would have made the locality particularly attractive to hominins in the past—something that is evidenced by the extraordinary accumulation of archaeological material recovered during survey of the area.

Survey of the basin resulted in the surface collection of artefacts along a number of transects across the tufa and surrounding basalt. Intensive survey using 5 × 5m grid squares was also carried out at site L0106, where a dense lithic scatter was discovered extending over about 100m<sup>2</sup> of the tufa surface, near to an area where the basalt outcrops through the tufa. Over 900 artefacts were collected from the survey area across a 40 × 50m area, which represents approximately a quarter of the total extent of this scatter. In total, 1002 lithic artefacts were recovered from within the Wadi Dabsa basin, including the surrounding basalt outcrops and the tufa. These display predominantly Early Stone Age/Lower Palaeolithic and Middle Stone Age/Middle Palaeolithic affinities, although several Later Stone Age artefacts produced exclusively from quartz were also found along the southern edge. The assemblage primarily consists of flake debitage, but also includes a large number of cores and several retouched tools (Table 1). Wadi Dabsa is the most artefact-rich location found thus far. Here we provide an initial analysis of the Acheulean material and its importance for elucidating early hominin landscape use within the Arabian Peninsula.





Figure 1. Location of Wadi Dabsa on the south-western coastline of the Arabian Peninsula.

## The Acheulean assemblage of Wadi Dabsa

A number of cores and retouched tools within the assemblage are typical of the Acheulean, including discoidal and simple flake cores with episodes of parallel working, as well as bifaces and large cutting tools. Nineteen of the artefacts can be classified as handaxes, cleavers or fragments thereof. Most of these tools were produced on large flakes, sourced either by deliberate flaking from large cores, or by selection of local, naturally produced exfoliation flakes. This method of production shares close similarities to other Acheulean sites within the Arabian Peninsula (Petraglia *et al.* 2009; Shipton *et al.* 2014), although the majority of the tools illustrate an intense focus on reduction of the tip, rather than the butt. High-quality basalts—almost certainly sourced from the surrounding lava fields—appear to be the predominant raw material of choice, with andesite used in much lower quantities. The local basalt from the lava fields, however, appears to vary in its porosity and density, with finer-grained materials to the north, and poorer-quality material along the southern edge (Inglis *et al.* 2015). The predominance of higher-quality raw materials within the assemblage, therefore, appears to indicate that the hominins present at the site carefully selected the better materials available.

**Table 1. Distribution of artefact types within the Wadi Dabsa assemblage.**

Type		Number found
<b>Flakes/debitage</b>	flakes	475
	prepared core flakes	96
	blades	17
	used flakes	28
	splintered pieces/wedges	3
	shatter	89
<b>Cores</b>	cores	140
	core fragments	6
<b>Bifacial tools</b>	handaxes	11
	cleavers	4
	pics	4
	broken handaxes	4
<b>Retouched tools</b>	backed knife	1
	burins	2
	denticulate	4
	notch	9
	large cutting tools	16
	piercers/borers	13
	points	8
	scrapers	47
<b>Other</b>	clasts	23
	hammerstones	2
	<b>Total</b>	<b>1002</b>



*Figure 2. Wadi Dabsa and associated geology and archaeological transects. L0107 (red highlight) indicates the location where the large handaxe was found.*




Figure 3. Top) photograph of the large handaxe from Wadi Dabsa; bottom) illustration of the handaxe, including profile view. Photograph by A. Shuttleworth; illustration by F. Foulds.

Within the assemblage, however, a single large bifacially worked tool stands out as anomalous (Figure 3). This was recovered during surface collection along a 250m transect at L0107, stretching from the north-western edge of the tufa to the top of a basalt jebel that overlooks the basin and wadi. It is 265mm long, weighs 3598g and was produced from either a very large basalt flake or, more probably, a natural exfoliation flake. On the basis of its size, it was originally interpreted as a large, abandoned roughout or core. Its appearance

**Table 2.** A comparison of the large handaxe with other known handaxes of length greater than 240mm. Data based on Leakey and Roe (1994) and Gowlett (2013).

Locality	Length (mm)	Breadth (mm)	Thickness (mm)
Kilombe	248	120	53
Kilombe	258	151	56
Kilombe	243	111	55
Sidi Abderrahman Cunette	250	162	47
Sidi Abderrahman Cunette	241	107	73
Kalambo Falls	291	138	65
Cornelia-Uitzoek	240	124	73
Cornelia-Uitzoek	243	114	77
Holsdam	245	107	65
Peninj	265	119	81
Olduvai Gorge FLK	289	132	72
Olduvai Gorge FLK	268	124	83
Olduvai Gorge FLK	249	116	72
Olduvai Gorge FLK	277	129	69
Olduvai Gorge FLK	270	117	67
<b>Wadi Dabsa</b>	<b>265</b>	<b>160</b>	<b>85</b>

shares affinities with Victoria West cores and with examples of cores developed on bifacial tools, albeit of a much larger size (DeBono & Goren-Inbar 2001; Sharon & Beaumont 2006; Sharon 2007, 2009). Limited preparation of the ventral surface and a lack of any additional examples from the site, however, preclude this interpretation. Furthermore, the large scar on the ventral surface appears to be a natural exfoliation surface, rather than an intentional removal. Evidence of bifacial retouch on the upper two-thirds using a heavy, hard hammer, and extensive working of the tip, probably using a smaller hard hammer, indicate the imposition of a working edge. This suggests that the artefact should be considered a finished tool, as opposed to an abandoned roughout, especially given that the pattern of reduction is closely comparable to similar examples of tip preparation seen on other bifaces recovered from the site.

Metrical analysis of large cutting tools (e.g. Sharon 2007) indicates that the large biface from Wadi Dabsa is well above average in terms of its size, even if it is not the largest currently known. A number of bifaces measuring at least 250mm have been found in both Europe and Africa, most notably those from Cuxton, Olorgesailie, Olduvai Gorge (site FLK), Isimila and the Furze Platt giant, all of which provide examples surpassing 300mm (Issac 1977: 134; MacRae 1987; Roe 1994: 207; Wenban-Smith 2004; Cole *et al.* 2016). A comparison of the Wadi Dabsa handaxe with several of these known large handaxes (Table 2) demonstrates that this new example fits well within the range of these previously collected artefacts, although it is generally broader and thicker than most. While the size of the large handaxe from Wadi Dabsa is comparable to others, it is rare that such tools approach weights of 3000g or more, with only a few known examples from Africa (Kelley 1959; Sharon 2007; Petraglia & Shipton 2008). The excessive weight of the example from




Figure 4. View from the top of a basalt jebel at the northern extent of L0107, looking south over the basalt surface and tufa exposure. Adapted from Inglis et al. (2015). Photograph by R. Inglis.

Wadi Dabsa would certainly have made it difficult to wield in the hand, begging the question of how and for what purpose this tool may have been used.

## Discussion

In the context of the wider Acheulean occupation of the Arabian Peninsula, Wadi Dabsa is comparable to sites such as Wadi Fatima and Dawādmī to the north, and those recently discovered in the Nefud Desert (Petraglia *et al.* 2009; Shipton *et al.* 2014; Jennings *et al.* 2015). It can also be added to the wider evidence for the Acheulean occupation of the Red Sea region produced by the DISPERSE Project and previous studies (Zarins *et al.* 1980, 1981; Inglis *et al.* 2013, 2014a & b, 2015). The location of Wadi Dabsa at the confluence of several tributaries and the potential presence of a larger body of water conform to the expectation that Acheulean sites are associated with water sources (Potts *et al.* 1999; Shipton 2011). This is unsurprising, given that hominin ranges would have been constrained by access to fresh water (Hardaker 2011). The surrounding basalt jebels would have provided expansive views of the surrounding landscape extending as far as the Red Sea coastline (Figure 4), which are equivalent to viewsheds reported for Wadi Fatima and Dawādmī (Petraglia *et al.* 2009).

The presence of large cutting tools (such as handaxes) produced on large flakes also conforms to descriptions of other Acheulean assemblages within Arabia. Although these cutting tools were produced using the abundant local raw materials, Wadi Dabsa displays clear evidence for the preferential selection of good-quality stone—specifically basalt clasts sourced to the north of the basin, which display a more cohesive cryptocrystalline structure compared to that available along the southern edge. This provides some evidence that

**Table 3.** A comparison of the mean length, thickness and weight of the Wadi Dabsa handaxes, with examples from Europe, Africa, India and the Arabian Peninsula. (\* Figures in brackets provide the average and standard deviations for the Wadi Dabsa assemblage following removal of the large handaxe.) Data based on Petraglia *et al.* (2009) and Shipton *et al.* (2014).

Locality	Mean length		Mean thickness			Mean weight		
	<i>n</i>	(mm)	<i>n</i>	(mm)	SD	<i>n</i>	(g)	SD
<i>Africa</i>								
Olduvai Gorge Bed II	21	195.39	17	66.92	19.2	17	1406.81	784.12
Kariandusi	58	157.94	35	43.6	14.74	35	571.02	369.8
Olorgesailie DE89A	63	180.76	60	46.23	10.43	60	877.82	381.8
Olorgesailie H9AM	13	199.77	10	36.2	7.53	10	770	426.54
Olorgesailie I3	62	97.95	57	33.54	9.28	57	225.12	197.48
Olorgesailie FB	16	98.81	15	34.6	8.44	15	180.87	116.11
Olorgesailie DE89C	69	158.7						
<i>Europe</i>								
High Lodge	68	116.51	63	35.15	14.01	63	259.89	208.83
<i>Arabia</i>								
Dawādmī 206–76	49	162.87	27	52.04	22.02			
Wadi Fatima	35	141.86	15	49.67	9.8			
Arzraq Lion Spring			42	43.97	9.68	42	216.43	86.11
<b>Wadi Dabsa*</b>	<b>11</b>	<b>140.27</b> <b>(127.8)</b>	<b>11</b>	<b>60.54</b> <b>(58.1)</b>	<b>15.83</b> <b>(14.33)</b>	<b>11</b>	<b>1105.72</b> <b>(856.5)</b>	<b>993.39</b> <b>(580.77)</b>
<i>India</i>								
Hunsgi V	151	143.51	45	48.44	9.99	45	669	349.6
Hunsgi II	34	162.9	18	52.22	10.6	18	1041.94	551.14
Gulbal II	17	147.14	12	47.5	9.65	12	902.5	385.84
Mudnur VIII	9	227.78	9	61.11	9.28	9	1302.22	204.56
Yediyapur I	21	123.13	10	36	5.16	10	443	230.3
Yediyapur IV	20	132.94	11	42.73	11.04	11	626.82	415
Yediyapur VI	66	127.86	21	42.86	13.09	21	591.19	563.49
Fatehpur V	31	126.82	11	40.91	11.36	11	455.45	246.74
Teggihalli II	31	121.54						
Anagwadi	25	137.24	15	45.73	6.04			
Godavari	10	114						

the local hominins had a clear appreciation of the variability in the conchoidal fracture properties of the lithic materials available to them. The presence of a Large Flake Acheulean at Wadi Dabsa, which is close to other Near and Middle Eastern sites that have been linked to similar knapping strategies seen at, for example, Gesher Benot Ya'aqov, suggests that these represent a new wave of Acheulean-using hominins dispersing from Africa (Martínez-Navaro & Rabinovich 2011). If this is the case, then Wadi Dabsa could expand this hypothesis to include the Arabian Peninsula.

The size and weight of the Wadi Dabsa handaxes fall within the range of variation generally recorded for the Acheulean (Table 3). In terms of shape, however, the handaxes found at Wadi Dabsa, including the large handaxe described above, show clear and repeated focus on reduction and finishing of the tip, leaving the butt minimally worked. This

suggests an active selection of a particular handaxe form. Variability in biface shape has long been a central topic within Lower Palaeolithic research. It has been suggested that variation in the shape of bifaces can often be explained by the need to establish and preserve a sharp cutting edge (Lycett 2008). A suite of factors, however, continues to be acknowledged as influencing handaxe shape, including raw material selection, social pressures and the individual (e.g. Ashton & McNabb 1994; Callow 1994; Gamble 1997; White 1998; Kohn & Mithen 1999; Spikins 2012; Foulds 2014). Among the bifaces from Wadi Dabsa, as well as the lithic artefacts from other sites examined as part of the DISPERSE Project, an emphasis on the creation of a good working edge is notable. It remains to be seen whether this was due to functional requirements, raw material affordance or the cultural transmission of specific methods of lithic manufacture in general.

The large handaxe presented here currently represents a unique find within the Arabian Peninsula, and is the largest handaxe from this region currently known to the authors. It falls within the range of variation seen amongst other examples of overly large tools, despite its excessive weight. The occurrence of only a single large biface at Wadi Dabsa, however, is more in keeping with the context in which such bifaces have been discovered in Europe, where they are generally found as single occurrences. That large handaxes are generally found in isolation, however, may present a false indication of their individuality. It is clear from African sites, where such large tools are found in assemblages (e.g. at Olduvai and Isimila—Roe 1994; Cole *et al.* 2016), that multiple, similar examples can occur. This may also be the case at Cuxton, where at least four handaxes over 200mm in length were recovered by Tester (Shaw & White 2003; Cole 2011), complementing the two large bifaces found during excavation by Wenban-Smith (2004).

The key question regarding the large handaxe is why it was produced. Several functional explanations have been posited on the phenomenon of large bifaces, including their role as digging tools, as expressions of knapping skill and as artefacts incorporated into some form of social display (Wymer 1968: 225, 1982: 103; Kohn & Mithen 1999). None of these theories has been convincingly proven. The large handaxe from Wadi Dabsa does not appear to represent the work of a highly skilled knapper wishing to demonstrate the extent of their abilities, whereas those used to support this hypothesis tend to be exquisitely worked (Wenban-Smith 2004). Prime examples are the biface from Furze Platt and the ficron and cleaver from Cuxton, which exhibit careful and controlled knapping to create a relatively well-thinned and symmetrical edge.

The excessive size and weight of the Wadi Dabsa biface leads us to believe that it was too large and unwieldy to be used in the hand—an observation that has been made of similar large tools (Wymer 1968, 1982; Roe 1981). Furthermore, it is unlikely that the maker intended to carry it from site to site. This suggests that either its use as a hand-held butchery tool, as is often proposed for handaxes, was unlikely, or alternatively, that our impressions of size and weight are significantly different to those of the hominins who made them (Wenban-Smith 2004). Conversely, it might be a large, bifacial core. As discussed above, however, the lack of additional examples and limited preparation appear to preclude this hypothesis. Despite the lack of extensive reduction used to form its overall shape, it seems reasonable to suggest that this large handaxe was made for a clear utilitarian purpose. This is supported by the fact that it conforms closely to other handaxes within

*A large handaxe from Wadi Dabsa and early hominin adaptations within the Arabian Peninsula*

the assemblage, most notably in the increased reduction intensity around the tip to create a cutting edge. It may perhaps have been employed as a static tool, with hominins resting the handaxe on the ground, secured between an individual's legs, and resources brought down on the tip for processing. In this way it could have been used to process faunal remains to access meat and marrow. Sites such as Isimila, Elandsfontein and Doornlaagte have provided examples of similar tools that were found on their edges when excavated, as if pressed into the ground (Wymer 1982: 103). While this is certainly plausible for the Wadi Dabsa handaxe, its recovery as part of an unstratified surface collection find from within the basalt fields means that this possibility cannot be substantiated. Microwear analysis of the tip will be required to determine whether it was used for a specific material or in a particular fashion.

## Conclusion

Wadi Dabsa presents a highly concentrated area of Acheulean activity within the Arabian Peninsula. It provided a wide range of resources, including raw materials for tool production and a fresh water source that would have attracted animals suitable for hunting. These resources were essential for hominin dispersal from the Red Sea shoreline and deeper into the Arabian Peninsula. The site is made more extraordinary by the large quantity of artefacts recovered, suggesting either the repeated or intensive use of this locality. The large handaxe adds to the complexity and difficulty of interpreting this newly discovered site, as well as representing a significant addition to the known catalogue of these enigmatic bifacial tools. While it is geographically unique, being the only example currently known from within the Arabian Peninsula, its unusually excessive weight highlights its unique nature in comparison to similar overly large tools. The use of such large bifaces is still a mystery, and it is hoped that the Wadi Dabsa specimen can contribute to this debate, as well as furthering discussion regarding their dispersal throughout the Acheulean world.

## Acknowledgements

We thank HRH Prince Sultan bin Salman bin Abdul Aziz, President of the Saudi Commission for Tourism and National Heritage (SCTH), Kingdom of Saudi Arabia; Ali Al-Ghabban, Vice-President; and Abdullah Al Saud, Director General, for granting fieldwork permission, as well as their interest in, and support of, our work in Saudi Arabia. Grateful thanks are also extended to Abdullah Al Zahrani, SCTH Riyadh, Saeed Al Karni, Director of Antiquities in Asir, and Haider Al Mudeer, as well as to the staff of the SCTH offices in Abha and Sabiya.

The January 2015 fieldwork was carried out as part of the DISPERSE Project, funded by European Research Council Advanced Grant 269586, under the 'Ideas' Specific Programme of FP7 to Geoff Bailey and Geoffrey C.P. King. For generous additional funding for the fieldwork, we thank the Gerald Averay Wainwright Fund for Near Eastern Archaeology, University of Oxford, and the Department of Archaeology Research Fund, University of York. This is DISPERSE contribution number 37.

## References

- ASEFAW, B., Y. BEYENE, G. SUWA, R.C. WALTER, T.D. WHITE, G. WOLDEGABRIEL & T. YEMANE. 1992. The earliest Acheulean from Konso-Gardula. *Nature* 360: 732–35.  
<https://doi.org/10.1038/360732a0>
- ASHTON, N. & J. McNABB. 1994. Bifaces in perspective, in N. Ashton & A. David (ed.) *Stories in stone* (Lithic Studies Society Occasional Papers 4): 182–91.

- BAILEY, G.N. 2009. The Red Sea, coastal landscapes and hominin dispersals, in M.D. Petraglia & J.I. Rose (ed.) *The evolution of human populations in Arabia: paleoenvironments, prehistory and genetics*: 15–37. Dordrecht: Springer.
- BAILEY, G.N., N. FLEMMING, G.C.P. KING, K. LAMBECK, G. MOMBER, L. MORAN, A. AL-SHAREKH & C. VITA-FINZI. 2007. Coastlines, submerged landscapes and human evolution: the Red Sea Basin and the Farasan Islands. *Journal of Island and Coastal Archaeology* 2: 127–60. <https://doi.org/10.1080/15564890701623449>
- BAILEY, G.N., G.C.P. KING, M. DEVÈS, N. HAUSMANN, R. INGLIS, E. LAURIE, M. MEREDITH-WILLIAMS, G. MOMBER, I. WINDER, A. ALSHAREKH & D. SAKELLARIOU. 2012. DISPERSE: dynamic landscapes, coastal environments and human dispersals. *Antiquity* 86(334) Project Gallery. Available at: <http://antiquity.ac.uk/projgall/bailey334/> (accessed 11 July 2017).
- BAILEY, G.N., M.H. DEVÈS, R.H. INGLIS, M.G. MEREDITH-WILLIAMS, G. MOMBER, D. SAKELLARIOU, A.G.M. SINCLAIR, G. ROUSAKIS, S. AL GHAMDI & A.M. ALSHAREKH. 2015. Blue Arabia: Palaeolithic and underwater survey in SW Arabia and the role of coasts in Pleistocene dispersals. *Quaternary International* 382: 42–57. <https://doi.org/10.1016/j.quaint.2015.01.002>
- CABRERA, V.M., K.K. ABU-AMERO, J.M. LARRUGA & A.M. GONZÁLEZ. 2009. The Arabian Peninsula: gate for human migrations out of Africa or cul-de-sac? A mitochondrial DNA phylogeographic perspective, in M.D. Petraglia & J.I. Rose (ed.) *The evolution of human populations in Arabia: paleoenvironments, prehistory and genetics*: 79–88. Dordrecht: Springer.
- CALLOW, P. 1994. The Olduvai bifaces: technology and raw materials, in M.D. Leakey & D.A. Roe (ed.) *Olduvai Gorge, volume 5: excavations in Beds III, IV and the Masek Beds, 1968–1971*: 235–53. Cambridge: Cambridge University Press.
- COLE, J. 2011. Hominin cognitive and behavioural complexity in the Pleistocene: assessment through identity, intentionality and visual display. Unpublished PhD dissertation, University of Southampton.
- COLE, J., J. McNABB, P. BUSHOZI, M. BATES, A. KWEKASON, D. NASH & P. TOMS. 2016. Recent investigations in the Stone Age site of Ismila, Tanzania. Paper presented at the Unravelling the Palaeolithic conference, Southampton, UK, 14–16 January 2016.
- DABBAGH, A., R. EMMERMAN, H. HÖTZL, A.R. JADO, H.J. LIPPOLT, W. KOLLMAN, H. MOSER, W. RAUERT & J.G. ZÖTL. 1984. The development of the Tihamat Asir during the Quaternary, in A.R. Jado & J.G. Zötl (ed.) *Quaternary Period in Saudi Arabia, volume 2: sedimentological, hydrogeological, hydrochemical, geomorphological, geochronological and climatological investigations in western Saudi Arabia*: 150–73. Vienna: Springer.
- DEBONO, H. & N. GOREN-INBAR. 2001. Note on a link between Acheulian handaxes and the Levallois method. *Proceedings of the Israel Prehistoric Society* 31: 9–23.
- DEVÈS, M.H., D.A. STURDY, G.C.P. KING, N. GODET & G.N. BAILEY. 2014. Hominin reactions to herbivore distribution in the Lower Palaeolithic of the southern Levant. *Quaternary Science Reviews* 96: 140–60. <https://doi.org/10.1016/j.quascirev.2014.04.017>
- FIELD, J.S. & M.M. LAHR. 2005. Assessment of the southern dispersal: GIS-based analyses of potential routes at Oxygen Isotopic Stage 4. *Journal of World Prehistory* 19: 1–45. <https://doi.org/10.1007/s10963-005-9000-6>
- FOULDS, F.W.F. 2014. Invisible individuals, visible groups: on the evidence for individuals and groups at the Lower Palaeolithic site of Caddington, Bedfordshire, UK, in F.W.F. Foulds, H.C. Drinkall, A.R. Perri, D.T.G. Clinnick & J.W.P. Walker (ed.) *Wild things: recent advances in Palaeolithic and Mesolithic research*: 12–40. Oxford: Oxbow.
- GAMBLE, C. 1997. Handaxes and Palaeolithic individuals, in N. Ashton, F. Healey & P. Pettitt (ed.) *Stone Age archaeology: essays in honour of John Wymer*: 105–109. Oxford: Oxbow.
- GOWLETT, J.A.J. 2013. Elongation as a factor in artefacts of humans and other animals: an Acheulean example in comparative context. *Philosophical Transactions of the Royal Society B* 368: article no. 20130114. <https://doi.org/10.1098/rstb.2013.0114>
- GROUCUTT, H.S. & M.D. PETRAGLIA. 2012. The prehistory of the Arabian Peninsula: deserts, dispersals, and demography. *Evolutionary Anthropology* 21: 113–25. <https://doi.org/10.1002/evan.21308>
- HARDAKER, T.R. 2011. *New approaches to the study of surface Palaeolithic artefacts: a pilot project at Zebra River, western Namibia* (British Archaeological Reports international series 2270). Oxford: Archaeopress.

- INGLIS, R.H., A.G.M. SINCLAIR, A. SHUTTLEWORTH & A.M. ALSHAREKH. 2013. Preliminary report on 2013 fieldwork in southwest Saudi Arabia by the DISPERSE Project: (2) Jizan and Asir provinces Feb–March. Unpublished report to the Saudi Commission for Tourism and Antiquities. Available at: <http://www.disperse-project.org/field-reports> (accessed 11 July 2017).
- INGLIS, R.H., A.G.M. SINCLAIR, A. SHUTTLEWORTH, N. HAUSMANN, A. AL MAAMARY, W. BUDD, M.G. MEREDITH-WILLIAMS, A.M. ALSHAREKH, S. AL GHAMDI & G.N. BAILEY. 2014a. Preliminary report on 2014 fieldwork in southwest Saudi Arabia by the DISPERSE Project: (1) Jizan and Asir provinces. Unpublished report to the Saudi Commission for Tourism and Antiquities. Available at <http://www.disperse-project.org/field-reports> (accessed 11 July 2017).
- INGLIS, R.H., A.G.M. SINCLAIR, A. SHUTTLEWORTH, A. ALSHAREKH, S. AL GHAMDI, M. DEVÈS, M.G. MEREDITH-WILLIAMS & G.N. BAILEY. 2014b. Investigating the Palaeolithic landscapes and archaeology of the Jizan and Asir regions, southwest Saudi Arabia. *Proceedings of the Seminar for Arabian Studies* 44: 193–212.
- INGLIS, R.H., F. FOULDS, A. SHUTTLEWORTH, A. ALSHAREKH, S. AL GHAMDI, A.G.M. SINCLAIR & G.N. BAILEY. 2015. The Palaeolithic occupation of the Harrat Al Birk: preliminary report on 2015 fieldwork in Asir province, southwest Saudi Arabia. Unpublished report to the Saudi Commission for Tourism and Antiquities. Available at <http://www.disperse-project.org/field-reports> (accessed 11 July 2017).
- ISAAC, G.L. 1977. *Olorgesailie: archaeological studies of a Middle Pleistocene lake basin in Kenya*. Chicago (IL): University of Chicago Press.
- JENNINGS, R.P., C. SHIPTON, P. BREEZE, P. CUTHBERTSON, M. ANTONIO BERNAL, W.M.C. OSHAN WEDAGE, N.A. DRAKE, T.S. WHITE, H.S. GROUCUTT, A. PARTON, L. CLARK-BALZAN, C. STIMPSON, A.-A. AL OMARI, A. ALSHAREKH & M.D. PETRAGLIA. 2015. Multi-scale Acheulean landscape survey in the Arabian Desert. *Quaternary International* 382: 58–81. <https://doi.org/10.1016/j.quaint.2015.01.028>
- KELLEY, H. 1959. Bifaces de tres grande taille. *Bulletin de la Société préhistorique française* 16: 739–72.
- KING, G.C.P. & G.N. BAILEY. 2006. Tectonics and human evolution. *Antiquity* 80: 265–86. <https://doi.org/10.1017/S0003598X00093613>
- KOHN, M. & S. MITHEN. 1999. Handaxes: products of sexual selection? *Antiquity* 73: 518–26. <https://doi.org/10.1017/S0003598X00065078>
- KÜBLER, S., P. OWENGA, S. RUCINA, S.J. REYNOLDS, G.N. BAILEY & G.C.P. KING. 2016. Edaphic and topographic constraints on exploitation of the Central Kenya Rift by large mammals and early hominins. *Open Quaternary* 2: article 5. <https://doi.org/10.5334/oq.21>
- LAMBECK, K., A. PURCELL, N.C. FLEMMING, C. VITA-FINZI, A.M. ALSHAREKH & G.N. BAILEY. 2011. Sea level and shoreline reconstructions for the Red Sea: isostatic and tectonic considerations and implications for hominin migration out of Africa. *Quaternary Science Reviews* 30: 3542–74. <https://doi.org/10.1016/j.quascirev.2011.08.008>
- LEAKEY, M.D. & D.A. ROE. 1994. *Oldovai Gorge, volume 5: excavations in Beds III, IV and the Masek Beds, 1968–1971*. Cambridge: Cambridge University Press.
- LEPRE, C.J., H. ROCHE, D.V. KENT, S. HARDMAN, R.L. QUINN, J.-P. BRUGAL, A. LENOBLE & C.S. FEIBEL. 2011. An earlier origin of the Acheulian. *Nature* 477: 82–85. <https://doi.org/10.1038/nature10372>
- LYCETT, S.J. 2008. Acheulean variation and selection: does handaxe symmetry fit neutral expectations? *Journal of Archaeological Science* 35: 2640–48. <https://doi.org/10.1016/j.jas.2008.05.002>
- MACRAE, R.J. 1987. The great giant handaxe stakes. *Lithics* 8: 15–17.
- MARTÍNEZ-NAVARRO, B. & R. RABINOVICH. 2011. The fossil Bovidae (Artiodactyla, Mammalia) from Gesher Benot Ya'aqov, Israel: out of Africa during the Early–Middle Pleistocene transition. *Journal of Human Evolution* 60: 375–86. <https://doi.org/10.1016/j.jhev.2010.03.012>
- PETRAGLIA, M.D. 2003. The Lower Palaeolithic of the Arabian Peninsula: occupations, adaptations and dispersals. *Antiquity* 77: 671–84. <https://doi.org/10.1017/S0003598X00061639>
- PETRAGLIA, M.D. & J.I. ROSE (ed.). 2009. *The evolution of human populations in Arabia: paleoenvironments, prehistory and genetics*. Dordrecht: Springer.
- PETRAGLIA, M.D. & C. SHIPTON. 2008. Large cutting tool variation west and east of the Movius Line. *Journal of Human Evolution* 55: 962–66.
- PETRAGLIA, M.D., N. DRAKE & A.M. ALSHAREKH. 2009. Acheulean landscapes and large cutting tool assemblages in the Arabian Peninsula, in M.D. Petraglia & J.I. Rose (ed.) *The evolution of human populations in Arabia: paleoenvironments, prehistory and genetics*: 103–16. Dordrecht: Springer.
- POTTS, R., A.K. BEHRENSMEYER & P. DITCHFIELD. 1999. Paleolandscape variation and Early Pleistocene hominid activities: members 1 and 7, Olorgesailie Formation, Kenya. *Journal of Human Evolution* 37: 747–88. <https://doi.org/10.1006/jhev.1999.0344>

- PRINZ, W. 1984. Geological map of the Wadi Haliy quadrangle, sheet 18E, Kingdom of Saudi Arabia. Scale 1:250 000. Geoscience Map GM-74C. Kingdom of Saudi Arabia Ministry of Petroleum and Mineral Resources, Deputy Ministry for Mineral Resources, Jeddah, Saudi Arabia.
- QUADE, J., N. LEVIN, S. SEMAW, D. STOUT, P. RENNE, M.J. ROGERS & S. SIMPSON. 2004. Palaeoenvironments of the earliest stone tool-makers, Gona, Ethiopia. *Geological Society of America Bulletin* 116: 1529–44. <https://doi.org/10.1130/B25358.1>
- REYNOLDS, S.C., G.N. BAILEY & G.C.P. KING. 2011. Landscapes and their relation to hominin habitats: case studies from *Australopithecus* sites in eastern and southern Africa. *Journal of Human Evolution* 60: 281–98. <https://doi.org/10.1016/j.jhevol.2010.10.001>
- ROE, D. 1981. *The Lower and Middle Palaeolithic periods in Britain*. London: Routledge & Kegan Paul.
- 1994. A metrical analysis of selected sets of handaxes and cleavers from Olduvai Gorge, in M.D. Leakey & D.A. Roe (ed.) *Olduvai Gorge, volume 5: excavations in Beds III, IV and the Masek Beds, 1968–1971*: 146–235. Cambridge: Cambridge University Press.
- ROHLING, E.J., K.M. GRANT, A.P. ROBERTS & J.-C. LARRASOANA. 2013. Paleoclimate variability in the Mediterranean and Red Sea regions during the last 500,000 years. *Current Anthropology* 54(S8): S183–201. <https://doi.org/10.1086/673882>
- SHARON, G. 2007. *Acheulian large flake industries: technology, chronology, and significance* (British Archaeological Reports international series S1701). Oxford: Archaeopress.
- 2009. Acheulian giant-core technology. *Current Anthropology* 50: 335–67. <https://doi.org/10.1086/598849>
- SHARON, G. & P. BEAUMONT. 2006. Victoria West—a highly standardised prepared core technology, in N. Goren-Inbar & G. Sharon (ed.) *Axe age: Acheulian tool making from quarry to discard*: 181–99. London: Equinox.
- SHAW, A.D. & M.J. WHITE. 2003. Another look at the Cuxton handaxe assemblage. *Proceedings of the Prehistoric Society* 69: 305–14. <https://doi.org/10.1017/S0079497X00001365>
- SHIPTON, C. 2011. Taphonomy and behaviour at the Acheulean site of Kariandusi, Kenya. *African Archaeological Review* 28: 141–55. <https://doi.org/10.1007/s10437-011-9089-1>
- SHIPTON, C., A. PARTON, P. BREEZE, R. JENNINGS, H.S. GROUCUTT, T.S. WHITE, N. DRAKE, R. CRASSARD, A. ALSHAREKH & M.D. PETRAGLIA. 2014. Large Flake Acheulean in the Nefud Desert of northern Arabia. *PaleoAnthropology* 2014: 446–62.
- SIDDALL, M., E.J. ROHLING, A. ALMOGI-LABIN, C. HEMLEBEN, D. MEISCHNER, I. SCHMELTZER & D.A. SMEED. 2003. Sea-level fluctuations during the last glacial cycle. *Nature* 423: 853–58. <https://doi.org/10.1038/nature01690>
- SPIKINS, P. 2012. Goodwill hunting? Debates over the ‘meaning’ of handaxe form revisited. *World Archaeology* 44: 378–92. <https://doi.org/10.1080/00438243.2012.725889>
- WENBAN-SMITH, F. 2004. Handaxe typology and Lower Palaeolithic cultural development: ficrons, cleavers and two giant handaxes from Cuxton. *Lithics* 25: 11–21.
- WHITE, M.J. 1998. On the significance of Acheulian biface variability in southern Britain. *Proceedings of the Prehistoric Society* 64: 15–44. <https://doi.org/10.1017/S0079497X00002164>
- WINDER, I., G.C.P. KING, M. DEVÈS & G.N. BAILEY. 2013. Complex topography and human evolution: the missing link. *Antiquity* 87: 333–49. <https://doi.org/10.1017/S0003598X00048985>
- WINDER, I., M. DEVÈS, G.C.P. KING, G.N. BAILEY, R.H. INGLIS & M. MEREDITH-WILLIAMS. 2015. Dynamic landscapes and complex topography as agents in human evolution: the dispersals of the genus *Homo*. *Journal of Human Evolution* 87: 48–65. <https://doi.org/10.1016/j.jhevol.2015.07.002>
- WYMER, J. 1968. *Lower Palaeolithic archaeology in Britain, as represented by the Thames Valley*. London: John Barker.
- 1982. *The Palaeolithic age*. London: Croom Helm.
- WYNN, T. 1995. Handaxe enigmas. *World Archaeology* 27: 10–23. <https://doi.org/10.1080/00438243.1995.9980290>
- ZARINS, J., N.M. WHALEN, I. IBRAHIM, A. MORAD & M. KHAN. 1980. Comprehensive archaeological survey program: preliminary report on the central southwestern provinces survey. *Atlat* 4: 9–117.
- ZARINS, J., A. MURAD & K. AL-YAISH. 1981. The second preliminary report on the southwestern province. *Atlat* 5: 9–42.

Received: 19 September 2016; Accepted: 22 December 2016; Revised: 17 January 2017